


## Anti-TurboGFP(d) antibody

Product	Cat.#	Lot.#	Size
Anti-TurboGFP(d) antibody	<b>AB513</b>	51301121268	100 µg

### Use

- Western blot
- Immunoblotting
- ICC
- ELISA



### Western blot detection of fluorescent proteins using anti-TurboGFP(d) antibody.

Lisates of HEK293 cells expressing fluorescent proteins were boiled in sample buffer (95 °C, 10 min) before loading. Anti-TurboGFP(d) antibody was used in the concentration 0.2 µg/ml. Secondary antibody: Goat anti-Rabbit HRP-conjugated IgG.

### Description

Rabbit polyclonal antibody against denatured and non-denatured TurboGFP and CopGFP.

**Specificity:** The antibody was selected to recognize denatured and non-denatured TurboGFP and CopGFP.

**Immunogen:** Full-length recombinant denatured TurboGFP and non-denatured CopGFP proteins comprising 6XHis tag.

**Antibody preparation:** Full-length recombinant denatured TurboGFP and non-denatured CopGFP proteins comprising 6XHis tag were purified from transformed *E. coli* using metal-ion affinity chromatography.

Antibodies were produced in rabbits immunized with the mixture of recombinant denatured TurboGFP and non-denatured CopGFP. Specific IgG were purified by TurboGFP affinity chromatography and mixed.

For affinity chromatography full-length recombinant TurboGFP was purified from transformed *E. coli* using organic extraction and ion exchange chromatography.

**Formulation:** Lyophilized from the PBS buffer containing 0.5% trehalose; pH 7.4.

**Reconstitution:** Reconstitute with sterile water or 50% glycerol to a concentration of 1 mg/ml.

**Storage:** Lyophilized samples are stable for twelve months from date of receipt when stored at -20 °C. The presence of silica gel drier is advisable.

Reconstituted with sterile water, antibody can be stored at 2 - 8 °C three months without detectable loss of activity.

Reconstituted with 50% glycerol, antibody can be stored at -20 °C in a manual defrost freezer for six months without detectable loss of activity. Aliquot antibody upon reconstitution. Avoid repeated freeze / thaw cycles.

### Recommendations for use

Anti-TurboGFP(d) antibody can be used to recognize denatured and non-denatured TurboGFP and CopGFP.

#### Working concentrations:

For Western blot use at a dilution of 1 : 1 000 - 1 : 5 000;

For ELISA use at a dilution of 1 : 30 000 - 1 : 50 000;

For immunocytochemistry use at a dilution of 1 : 1 000 - 1 : 3 000.

**Note.** Optimal dilutions/concentrations should be determined by the end user.

**Tissue (cells) fixation for immunohistochemistry:** Formaldehyde (formalin, paraform) fixation is recommended. For example, tissues can be fixed in PBS containing 4% formaldehyde for 10-15 min, treated with 0.1% saponin in PBS for 10-15 min, and washed three times in PBS.

**Sample preparation for Western blot:** To a sample containing 10-100 ng of a target protein, add an equal volume of 2X SDS-PAGE sample buffer. Heat the sample at 95 °C before loading on a gel or spotting on a membrane (for dots).

### Notice to Purchaser:

These products are intended for research purposes only.

**MSDS information** is available at <http://evrogen.com/support/MSDS-info.shtml>